

Report on  
**1:100 000 Scale Geological and Metallogenic Maps**  
**Sheet 3166-23**  
Province of Córdoba

Patrick Lyons and Roger G. Skirrow

*GEOSCIENTIFIC MAPPING OF THE SIERRAS PAMPEANAS ARGENTINA-AUSTRALIA  
COOPERATIVE PROJECT*

**AUSTRALIAN GEOLOGICAL SURVEY ORGANISATION**

1997

# CONTENTS

## SECTION 1: GEOLOGY

### 1. INTRODUCTION 1

- 1.1 Location and access 1*
- 1.2 Nature of work and previous investigations 1*

### 2. STRATIGRAPHY 3

- 2.1 General relations 3*
- 2.2 Early Palaeozoic metamorphic basement 5*
  - Marble (€l) 5
  - Pichanas Metamorphic Complex and associated S-type granites (€gnp, €gno, €gnt, €gnh, €eb, €gi1, €g) 6
- 2.3 Plutonic rocks 10*
  - Esmeralda Granodiorite (Dgr) 10
  - La Playa Aplo-pegmatite (Dgp1) 11
  - La Playa Granodiorite (Dgp2) 12
  - Granite, unassigned (Dg) 12
  - Minor dyke rocks 12
- 2.4 Cainozoic 13*
  - 2.4.1 Tertiary 13*
    - La Playa Formation (Tpp) 13
  - 2.4.2 Tertiary to Quaternary 13*
  - 2.4.3 Quaternary 14*

### 3. TECTONICS 14

- 3.1 Pampean Cycle: Early Cambrian deformation and metamorphism 15*
- 3.2 Famatinian Cycle: Ordovician deformation and metamorphism 16*
- 3.3 Achaian Cycle: Devonian deformation and metamorphism 17*
- 3.4 Andian Cycle: reverse faulting 17*

### 4. GEOLOGICAL HISTORY 18

- 4.1 Early Cambrian sedimentation 18*
- 4.2 Pampean Cycle 19*
- 4.3 Famatinian Cycle 19*
- 4.4 Achaian Cycle 20*
- 4.5 Carboniferous - Permian sedimentation 21*
- 4.6 Andean Cycle 22*

## **SECTION 2: ECONOMIC GEOLOGY**

### **1. INTRODUCTION 23**

### **2. METALLIC MINERAL OCCURRENCES 24**

### **3. NON-METALLIC MINERALS AND *ROCAS DE APLICACION* 24**

*3.1 Limestone, Marble 25*

*3.2 Granite 25*

*3.3 Fluorite, Clay, Ochre, Steatite, Garnet and Amphibolite 25*

*3.4 Mica, Quartz, Feldspar 25*

### **BIBLIOGRAPHY 26**

### **ARGMIN DATABASE OUTPUT SHEETS 30**

# SECTION 1: GEOLOGY

by Patrick Lyons

## 1. INTRODUCTION

### 1.1 LOCATION AND ACCESS

The 3166-23 1:100 000 Sheet area lies within Córdoba Province, between 31°00'-31°20'S and 65°00'-65°30'W. The area is part of the 3166-IV (Villa Dolores) 1:250 000 sheet area.

The region includes the western Sierra Grande which is drained by the north flowing Ríos Pichanas and Guasapampa.

Access to the region, from Córdoba city, is via Ruta Nacional 38 which connects the main population centres of La Cumbre, Capilla del Monte, Cruz del Eje, and Villa de Soto. A number of secondary roads connecting the centres of La Higuera, Ciénaga del Coro, and La Playa to the main centres afford good access to most of the rock types in the sheet area.

### 1.2 NATURE OF WORK AND PREVIOUS INVESTIGATIONS

Mapping of the Sierras septentrionales de Córdoba was carried out in 1995 and 1996 under the Geoscientific Mapping of the Sierras Pampeanas Argentina - Australia Cooperative Project by geologists from the Australian Geological Survey Organisation (AGSO) and the Subsecretaria de Minería (DNSG) (Figure 1.1). The mapping employed a multidisciplinary approach using the newly acquired high-resolution



**Figure 1.1.** Location and simplified geology of the Sierras septentrionales de Córdoba and location of 1:100 000 sheets.

airborne magnetic and gamma-ray spectrometric data, Landsat TM imagery, and 1:20 000 scale (approximate) black and white air photography. All geological maps were compiled on topographic bases produced at photo-scale from rectified Landsat images controlled by field GPS sites. Geologists involved in the fieldwork were P. Lyons, and P.G. Stuart-Smith (AGSO), and J.C. Canadiani, H. Lopez, and R. Miro (DNSG). P. Espejo, M. Viruel, D. Martos (DNSG) and B. Torres (Secretaría de Minería de la Provincia de Córdoba) assisted with the fieldwork.

The area was first mapped as undifferentiated metamorphic basement by early workers (e.g. Pastore, 1932). Much of the Sierra Grande was mapped by Olsacher between 1961 and 1964 (Lucero Michaut and Olsacher, 1981)

## **2. STRATIGRAPHY**

### **2.1 GENERAL RELATIONS**

The 3166-23 Sheet area is part of the southern Sierras Pampeanas, a distinct morphotectonic province of early to mid Palaeozoic metamorphic, felsic and mafic rocks, forming a series of block-tilted, north-south oriented ranges separated by intermontane basins (Figure 2.1). The ranges are bounded by escarpments developed on moderate to steeply dipping reverse faults developed during the Cainozoic Andean uplift (Jordan and Allmendinger, 1986).

Basement consists of Early Cambrian metamorphic and igneous complexes intruded by Cambrian, Ordovician, and Devonian granitoids. The remnants of an extensive Carboniferous-Permian fluvio-lacustrine sequence (González and Aceñolaza, 1972) occur in a graben valley in the north west of the map area and continental Cainozoic and Quaternary cover occupies major valleys and intramontane areas to the north, east and west.



**Figure 2.1.** Location of the three project areas of the Argentina-Australia Cooperative Project and simplified regional geology of the southern Sierras Pampeanas.

Recent geological and geophysical surveys conducted as part of the Cooperative Argentine-Australia project in the Sierras Pampeanas show that the Paleozoic basement of the southern Sierras Pampeanas contains of a number of distinct lithological and structural domains separated by major tectonic zones. There are two principal domains: a Cambrian Pampean domain, and the Ordovician Famatinian domain to the west. Both domains have shared a common geological history since early Ordovician times. The boundary between the domains is broadly coincident with a regional change in the gravity near the western flank of the Sierras de Córdoba (Miranda and Introcaso, 1996) and is marked by the Guzman Fault further south in the Province of San Luis. Only the Pampean domain is exposed in the Sierras septentrionales de Córdoba. However, the younger Famatinian domain is inferred to be present in the subsurface west of the Sierra Grande. A summary of stratigraphy and relations is given in Table 2.1.

## 2.2 EARLY PALAEOZOIC METAMORPHIC BASEMENT

### Marble (e1)

Minor marble units are found throughout the sheet area but mappable outcrop is principally found in the Pichanas Metamorphic Complex in the south of the sheet area near the Río Pichanas and Arroyo Ciénaga del Coro. Details are given under the respective sections.

**Table 2.1.** Summary of stratigraphy and relationships in the 3166-18 1:100 000 Sheet.

Age (Ma)	Unit	Description	Relations
----------	------	-------------	-----------



QUATERNARY	Talus deposits.	Unconsolidated debris of basement material	Deposits along the fault scarps Sierras Serrezuela and Guaspampa
	Alluvium	Unconsolidated clay, sand and gravel	Deposits along active river courses
CAINOZOIC	Undifferentiated fluvial and aeolian deposits	Clay, sand, gravel, paleosol	Mantles older units.
	Silcrete	Thin unit about 1-3m thick	Covers Miocene volcanoclastics.
TERTIARY	La Playa Formation	Travertine	
CARBO-PERMIAN	Tasa-Cuna Formation	Shale, sandstone, conglomerate	Remnant preserved in graben.
DEVONIAN	La Playa Aplopegmatite	Granite	Intrudes La Playa Granodiorite and Pichanas Formation.
	La Playa Granodiorite	Granodiorite	Intrudes Pichanas Complex.
	Esmeralda Granodiorite	Granodiorite	Intrudes Pichanas Complex.
	Unassigned granite	Leucogranite	Intrudes Pichanas Complex
CAMBRIAN	Pichanas Complex	Pelitic gneiss, quartzite, migmatite, S-type granite, minor marble and amphibolite.	Concordant(?) with Cruz del Eje Complex.

**Pichanas Metamorphic Complex and associated S-type granites (εgnp, εgno, εgnt, εgnh, εeb, εgi1, εg)**

*Paragneiss, quartzite, schist, migmatite, associated S-type granite, minor marble and amphibolite*

The Pichanas Metamorphic Complex is the most extensive unit in the sheet area. Its name derives from the Río Pichanas. It is primarily composed of cordierite and garnet paragneiss and migmatite with minor marble and amphibolite units. In places, high T metamorphism resulted in local melting which produced a number S-type granite

The Pichanas Metamorphic Complex shares a common deformational and metamorphic history with the Cruz del Eje Metamorphic Complex to the west (Sheets 3166-18,

3166-24) and a Th-Pb age determined from monazite gave  $526 \pm 11$  Ma as the age for peak amphibolite grade metamorphism (Camacho and Ireland, 1997).

Metapelitic paragneisses with minor marble and amphibolite ( $\epsilon_{gnp}$ ,  $\epsilon_{gno}$ ) form the dominant units in the complex. They grade into migmatite in places. They contain medium- to high-grade assemblages of quartz (25% to 40%), K-feldspar (25% to 35%), biotite (10% to 30%), plagioclase (5% to 10%), muscovite (5% to 10%) and minor garnet, cordierite, and sillimanite. Retrograde effects are evident by muscovite replacement of sillimanite and K-feldspar and chlorite replacement of cordierite and garnet. They are identified by the relative proportions of cordierite and garnet; cordierite is greater than garnet in  $\epsilon_{gnp}$  and garnet is greater than cordierite in  $\epsilon_{gno}$ . However, the boundary between the two is gradational. This may reflect bulk compositional differences or facies changes in the original sedimentary pile. On radiometric images a broad distinction can be made as the garnetiferous unit gives higher uranium and thorium counts.

A unit known as in earlier works as the Tuclame Formation ( $\epsilon_{gnt}$ ), e.g. Lucero Michaut and Olsacher (1981), crops out extensively south of Paso Viejo and as a narrow north-west trending belt, about 1 km to 2 km wide, extending from west of the Río Pichanas to the Tasa Cuna Valley.

Although structurally complex, it appears to be concordant with the cordierite dominant paragneiss except for the north east side of the aforementioned belt which is separated by a faulted informally called the Dos Pozos mylonite-cataclasite zone. The zone is about 100m wide, sub-vertical, and kinematic indicators show a north-east side up movement. Whether this is the only movement on the fault is not known. The fault does not offset Early-Mid Devonian mineralisation which transects it.

It comprises a sequence of non magnetic (magnetic susceptibilities  $< 20 \times 10^{-5}$  SI) *quartz-rich meta psammitic and minor pelitic gneiss* which are typically banded on a centimetre scale. The banding consists of alternating quartzo-feldspathic layers with thinner micaceous-rich layers. Major mineral constituents are quartz (40% to 70%), biotite (15% to 30%), muscovite (5% to 25%), plagioclase (10% to 25%), K-feldspar ( $< 30\%$ ),

and sillimanite (< 10%) which is mostly retrogressed to muscovite. Accessory phases include tourmaline, apatite, rutile, and zircon. The presence of migmatite pods, derived from partial melting of pelitic units within the original sedimentary pile, indicates that peak metamorphism was at least amphibolite facies. It is similar to rocks within the Cruz del Eje Formation and the Conlara Complex in an Luis Province and the Olta Complex in La Rioja Province. As a possible correlation may be made it has not been formally separated here.

This unit preserves a complex Pampean progressive deformation history, involving up to three folding events, which are not apparent in the gneisses and migmatites of other units in the Sierras septentrionales. In places, the banding is clearly a differentiated crenulation cleavage and in lower grade parts it may also represent original bedding.

*Quartz, K-feldspar, biotite, plagioclase,  $\pm$ chlorite,  $\pm$ sillimanite,  $\pm$ cordierite* gneiss ( $\in$ gnh) crops out over a small area extending from the north eastern margin of the La Playa Granodiorite to a point about 3 km north of Agua de Ramón. Due to poor access through the unit it is relatively unknown but may be distinguished by slightly elevated biotite contents compared to the cordierite and garnet gneisses to the east, and zones of granodioritic composition which may be orthogneiss. The unit has a distinct radiometric signature with elevated K and Th responses relative to adjacent units, which may be due to a concentration of pegmatite or the presence of metasomatism related to mylonitisation apparent on its western margin where kinematic indicators indicate east over west movement with a component of sinistral displacement. The regional gneissic foliation generally trends north-south but is locally east-west in the area about a kilometre north of Agua de Ramón.

*Quartz, muscovite, biotite,  $\pm$ K-feldspar,  $\pm$ cordierite,  $\pm$ garnet* schist ( $\in$ eb) crops out in the far north of the Sierra de Guasapampa and as small scattered exposures and inliers west of the escarpment. It covers an area of about 12 km<sup>2</sup>. Previously the schist was mapped as tonalitic biotite gneiss (Lucero Michaut and Olsacher, 1981). The indistinct contact with the gneiss to the south and may represent a slight change in grade or an original sedimentary facies variation.

The mineral assemblage is characteristic of medium to high grade pelitic schist, and has a muscovite content of about 30%. The schistosity, with common intrafolial folds is sub-horizontal to gently east dipping. A moderate to steeply east-dipping, crenulation cleavage is developed in places and steepens in the east near the Río de Guasapampa where it becomes more penetrative with a corresponding increase in metamorphic grade.

Bodies of S-type granite associated with Early Cambrian high grade metamorphism occur throughout the Pichanas Metamorphic Complex. The most well known of these, and its cogenetic cordierite, has been previously named the El Pilón Granite or El Pilón Formation (Lucero Michaut and Olsacher, 1981) as its distribution was formerly recognised only over a small area between La Represa and Los Simbolitos (3166-18 Sheet) a distance of about 6 km. However, it is the main body of a discontinuous belt about 20 km long, which crops out as small hills and turtle backs between Los Simbolitos and Embalse Pichanas Concordant and discordant contacts with the surrounding meta-pelites, and numerous enclaves of the same, provide field evidence that these granite bodies are voluminous accumulations of partial melt products generated during high-T metamorphism of the pelites.

On this sheet it K-feldspar phyric granite(ε<sub>g1</sub>) is the only assigned S-type granite. It is a pink to deep pink, *phyric to megaphyric, K-feldspar biotite granite* with minor muscovite, sillimanite, chlorite, after biotite, and cordierite and trace amounts of plagioclase and zircon. K-feldspar laths often display a local flow alignment, and make up about 40% to 45% of the granite. Biotite contents are about 15% to 20%. The radiometric response is high with total counts around 90 cps to 95 cps and potassium around 6.5 cps which gives a clearly identifiable response on radiometric images. Magnetic susceptibility is low, generally about  $10 \times 10^{-5}$  SI.

Recent isotopic age data obtained for the S-type granites are summarised in Table 2.2. Although recent U-Pb zircon data (Camacho and Ireland, 1997) are approximate, due to some Pb loss, there is broad agreement with Rb-Sr ages obtained by Rapela and others (1995).

**Table 2.2.** Isotopic ages of S-type granite from the Pichanas Metamorphic Complex. 1. Rapela and others (1995) 2. Camacho and Ireland (1997).

Rock type	Age (Ma)	Method	Ref.	Comment
Granite	520±5	Rb-Sr	1	Leucogranite (and cordierite rock)
Granite	ca 527	U-Pb	2	Phyric granite
Granite	ca 514	U-Pb	2	Leucogranite

Some lead loss at about 480 Ma (Camacho and Ireland, 1997) may be due to the Ordovician Famatinian event.

Small bodies of unassigned granite (εg), interpreted from aerial photographs and radiometric images, occur as part of a broad belt trending NW-SE from an area about 3 km east of Tosno to area about 8 km west of La Higuera and as smaller isolated pods around the main body.

Limited field evidence suggests they are porphyritic to equigranular, ranging in composition from granite to granodiorite. They contain numerous enclaves of schist and gneiss derived from meta-pelites and typically form intrusive, elongate to equidimensional, bodies which are concordant to sub-concordant with the regional fabric of the enclosing metamorphic rocks. They probably originated as melt products derived during high-T regional metamorphism of the metasediments. There are no isotopic age determinations, however, an Early Cambrian age is interpreted as the age of the peak regional metamorphism in the area is about 530 Ma (Camacho and Ireland, 1997). The main body, which has similar field relationships to country rock as the undifferentiated bodies, crystallised at about 520 Ma (Rapela and others, 1995; this report). These ages are consistent with an origin as anatectic melt products (cf. Fitzsimmons and others, 1996).

## 2.3 PLUTONIC ROCKS

### Esmeralda Granodiorite (Dgr)

The Esmeralda Granodiorite is a small body, covering about 3 km<sup>2</sup>, immediately east of Agua de Ramón. It is well exposed and easy access is afforded by disused roads connecting the W deposits which it hosts. It is named after the Mina Esmeralda which is the largest of these deposits.

It is equigranular, medium grained, weakly foliated, and contains quartz, K-feldspar, plagioclase, biotite, muscovite, both primary and secondary, and minor to trace sericite, chlorite, epidote, and zircon. Lapidus and Rossi (1959) report the presence of apatite. Limited whole rock geochemistry indicate it is a granodiorite, albeit that it has an elevated soil content which may be due to silicification related to mineralisation. The foliation is parallel to the steep NNE striking regional foliation. Magnetic susceptibility is low, less than  $10 \times 10^{-5}$  SI, and the radiometric response is weak. Magnetic images suggest the western margin of the granodiorite is fault bounded. There are no isotopic age data for the Esmeralda Granodiorite, however, there is a possible genetic relationship between the granodiorite and W mineralisation. The <sup>40</sup>Ar-<sup>39</sup>Ar dating of muscovites from cross cutting tungsten bearing veins gave ages between 360 Ma and 368 Ma (Camacho, 1997).

### **La Playa Aplo-pegmatite (Dgp1)**

This is a high K granite which occupies a small area on the east of the La Playa Granodiorite which it intrudes and appears to be genetically related to the prominent aplo-pegmatites found within it. Although a dominantly medium grained rock, it contains substantial patches of very coarse grained pegmatitic material. Modal analyses (R. Miró, pers. comm.) show it contains about 35% to 40% quartz, 30% to 40% plagioclase, 20% to 30% K-feldspar, 2% to 7% muscovite, and 2% to 6% biotite. Minor and trace minerals include garnet, epidote, apatite, rutile, and zircon. There are no isotopic data but the close spatial association and intrusive relationship with the La Playa Granodiorite suggest a Devonian age.

### **La Playa Granodiorite (Dgp2)**

This granite crops out immediately west of La Playa and covers an area of about 30 km<sup>2</sup>. It is a white to light grey, equigranular, medium grained, biotite bearing granodiorite which intruded rocks of the Pichanas complex and Totora Huasi Gneiss. Gomez (via R. Miró, pers. comm.) gives modal analyses of about 35% quartz, 35% plagioclase, 12% microcline, and 15% biotite, including secondary chlorite, and accessory muscovite, primary and secondary, epidote, sericite, and zircon. It is intruded by prominent aplo-pegmatite dykes genetically related to Dgap. There are no age data for the La Playa Granodiorite, but similarities between its major and trace element geochemistry and that of the Esmeralda Granodiorite and their spatial proximity suggests a Devonian age.

### **Granite, unassigned (Dg)**

A small body of leucogranite about 7 km south of Embalse Pichanas crops out as small bosses over an area of less than 5 Ha. It is a white, equigranular, medium grained granite similar in hand specimen to the La Playa Granodiorite (Dgp2) containing K-feldspar, quartz, plagioclase, muscovite, and minor biotite and chlorite. It also has similar magnetic and radiometric responses to the La Playa Granodiorite and is possibly an apophysis of a genetically related body. There are no geochemical or isotopic data but the above superficial similarities suggest it is the same age as the La Playa body.

### **Minor dyke rocks**

#### *Pegmatite*

Several generations of pegmatite dykes intrude basement metamorphics and granitic rocks. The oldest form small deformed pods, less than a metre wide. These pegmatites are probably the product of partial melting during Cambrian M1 metamorphism (Pampean).

Pegmatite dykes swarms are common in quartzite units (εgnt) of the Pichanas Formation.

Ridge forming pegmatites related to the La Playa Aplopegmatite intruded the La Playa Granodiorite.

### *Lamprophyre*

Lapidus and Rossi (1959) report west to north-west dipping lamprophyre dykes intruded into basement gneiss in the vicinity of the Mina El Carmen tungsten deposit near Agua de Ramón. They are cut by late Devonian tungsten bearing veins.

## 2.4 CAINOZOIC

### 2.4.1 Tertiary

#### **La Playa Formation (Tpp)**

##### *Travertine*

The Pliocene La Playa Formation (Olsacher, 1960) occurs as small out crops in the vicinity north and south of La Playa. Possibly related to Cainozoic volcanism (Lucero Michaut and Olsacher, 1981), it consists of light grey, fine-grained calcite with occasional layers of carbonaceous material and lithic fragments deposited in a lacustrine environment. Fossils of plant stems and fresh water molluscs are common.

Small cavities are lined with fine-grained crystalline quartz.

### 2.4.2 Tertiary to Quaternary



An extensive Cainozoic sequence (Czu) of *undifferentiated Tertiary to Quaternary fluvial and aeolian deposits and paleosols* which cover a large part of the Pampean region. The upper loess portion of the unit has been mapped in the area east of the Sierra Chica as the ?Upper Pleistocene “Formación General Paz” by Santa Cruz (1978). Strasser and others (1996) correlated similar deposits in the San Luis region with Late Pleistocene and Holocene units in the Buenos Aires Province.

The unit is dominated by pinkish loess which covers all older units forming a mantle or rarely dune fields which cover and preserve pre-existing topographic relief between the main Pampean ranges. Present river courses and associated deposits locally dissect the deposits.

The loess deposits comprise mostly friable illite and silt. Material was derived from both the metamorphic-igneous basement rocks and a volcanic-pyroclastic source (Santa Cruz, 1978).

### **2.4.3 Quaternary**

Holocene (Santa Cruz, 1978) to Recent *alluvial deposits of clay, sand and gravel* (Qa) occur along active river courses and adjacent terraces. The most extensive deposits are associated with the Ríos Soto and Pichanas in the north.

Minor Recent *talus deposits* (Qt) of granitic debris, derived from basement rocks, occur on the western slopes of the Sierra Serrezuela and Sierra Guasapampa. The steep slopes are all reverse fault scarps formed during the Cainozoic Andean Cycle 2.3

## **3. TECTONICS**

Three major deformation/metamorphic and magmatic events have affected basement rocks: the Early Cambrian Pampean Cycle, the Early Ordovician Famatinian Cycle and the Devonian Achalian Cycle. Faulting and block-tilting occurred during the Mesozoic and later Cainozoic Andean Cycle.

### 3.1 PAMPEAN CYCLE: EARLY CAMBRIAN DEFORMATION AND METAMORPHISM

No original sedimentary structures, such as bedding, are unequivocally recognised in the metamorphic rocks. In general, the earliest surface (S1) is a penetrative gneissic foliation in pelitic gneiss, defined by leucosome lenses and bands and foliated muscovite-biotite-rich bands. In amphibolite and calc-silicate rocks the foliation forms strongly differentiated mineralogical bands with aligned hornblende. Throughout most of the region the S1 foliation forms the dominant trends on aerial photographs and satellite imagery.

Sillimanite-garnet assemblages in pelitic gneiss indicate M1 metamorphism was at least upper amphibolite facies and abundant muscovite-pegmatites and leucosome (forming subconcordant lenses with S1), suggest limited partial melting took place. Estimates of peak metamorphic conditions for the Sierra Chica are mostly about 6 Kb, and 700°C to 800°C (see Table 3.1). Uranium-lead dating of zircon rims and monazite grains which grew during the peak metamorphic event (M1) in both the Pichanas and Cruz del Eje Metamorphic Complexes, give an age of about 530 Ma, interpreted here as the age of the M1/D1 event of the Pampean Cycle.

**Table 3.1.** M1 Metamorphism - estimated peak conditions in the southern Sierras Pampeanas

Location	Temperature (°C)	Pressure (Kb)	Reference
Sierras de Córdoba	700 - 750	6.1 - 6.4	Toselli & others, 1992
Sierra Chica	500 - 700	4 - 6	Pérez & others, 1996
Sierra del Cuniputo-Totoralejo	700 - 800	6 - 8	Murra & Baldo, 1996
Sierras Comechingones	700 - 750	6.1 - 6.4	Cordillo, 1984
Sierra Chica de Córdoba	820	5.7	Baldo & Casquet, 1996
Sierras Comechingones	760-800	8.5-9.5	Cerredo, 1996
San Carlos	650-700	4.5-5	Demange & others, 1993
Sierras Comechingones/ Chicas	800	8	Martino & others, 1994

In the Sierra Grande there are, broadly, two principal orientations of the D1 high grade fabric (local S3 or S4 ?), steeply dipping to the south west and north-north east. However, in the Sierra de Guasapampa the fabric is upright or dips steeply to the west. This fabric is tightly to isoclinally folded at meso-scale. Limited data show the plunge of meso-scale fold hinges unevenly distributed about a girdle oriented steeply to the north west. This suggests that the fabric was produced during regional ductile shearing, however, shear sense indicators are absent.

### 3.2 FAMATINIAN CYCLE: ORDOVICIAN DEFORMATION AND METAMORPHISM

Widespread isoclinal folding and thrusting, at lower amphibolite/upper greenschist facies conditions, throughout the region is attributed to the Early Ordovician Famatinian Cycle. Metamorphic conditions produced chlorite from garnet and cordierite, and muscovite from sillimanite.

In areas of penetrative S1 development all planar D1 fabric elements are rotated into parallelism with the S2 foliation with a pronounced mineral lineation (L2) of biotite, muscovite and quartz, which plunges shallowly to the east (~ 100°). This lineation is perpendicular to long axes of boudinaged tonalitic layers, indicating a broadly coaxial deformation.

The age of this deformation is poorly constrained in the Sierra Septentrionales de Córdoba. K-Ar and Rb-Sr dating of muscovite from a pegmatite which was emplaced syn D2 yields a minimum age of about 428 Ma (Camacho,1997). The deformation (D2) is interpreted here as part of the Early Ordovician Famatinian Cycle which is dated in the Sierras de San Luis at 490 Ma (Camacho. 1997).

Sims and others (1997) have identified a *compressional* and an *extensional* phase of the Famatinian event in Sierras de San Luis. However, no evidence for a widespread extensional regime has been found in the sierras septentrionales de Córdoba.

Boudinaged phyric granites in the Guamanes Shear Zone (Sheet 3166-18, 24) predate the final Devonian compression and may have intruded during the Famatinian extensional regime.

### **3.3 ACHALIAN CYCLE: DEVONIAN DEFORMATION AND METAMORPHISM**

A complex system of rectilinear brittle subvertical sinistral NW- and dextral NE-trending strike-slip faults, breccia zones, fractures and kink zones (S4) affect all the basement units in the Sierras Septentrionales de Córdoba, crosscutting the S3 foliation where present. Regional faults are rarely exposed, but are prominent on aerial photographs and Landsat images. Some of the faults are also delineated on magnetic images as zones of demagnetisation.

The orientation and conjugate relationship of the WNW- and NE-trending strike-slip faults, breccia zones and fractures indicates a possible continuation of the east-west compressive regime that accompanied S3 and S4 development. This fracture system is developed throughout the Sierras Pampeanas, and in Córdoba and La Rioja Provinces where muscovite Ar-Ar ages of micas from quartz veins indicate that this stage began about 385 Ma, peaked at 370 Ma and continued until 355 Ma (Camacho, 1997). These faults zones therefore represent the final stage of the Achalian Cycle.

### **3.4 ANDIAN CYCLE: REVERSE FAULTING**

The Sierras de Serrezuela and Guasapampa are examples of basement tilt blocks formed by east-west compression during the Cainozoic Andean uplift (Jordan and Allmendinger, 1986). The ranges slope gently to the east and are bounded to the west by escarpments developed on moderate to steep east-dipping reverse faults. Many of the faults are probably reactivated Palaeozoic or Mesozoic structures. Quaternary faulting effects are limited to characteristic haematitic zones of fault gouge up to 5 m wide which dip 30-55° to the east, crosscutting older and mostly steeper fault fabrics.

## 4. GEOLOGICAL HISTORY

The Sierras Septentrionales de Córdoba area form part of the southern Sierras Pampeanas, comprising basement ranges of early Palaeozoic metamorphic rocks and Palaeozoic granitoids, separated by intermontane Mesozoic and Cainozoic sediments. The basement rocks form a series of north-trending lithological and structural domains separated by major mid-crustal shear zones. These domains have been variously interpreted to form (originally) part of an ensialic mobile belt (e.g. Dalla Salda, 1987) or as terranes which either accreted or developed on a western convergent margin of the Río Plata craton (e.g. Ramos, 1988; Demange and others, 1993; Escayola and others, 1996, Kraemer and others, 1995, 1996). Recent geochronological studies indicate that there are two principal domains in the southern Sierras Pampeanas: an older Cambrian Pampean domain, and a younger Ordovician Famatinian domain to the west, not exposed in the map area. Both domains share a common geological history since early Ordovician times. The boundary between the domains is broadly coincident with a regional change in the gravity on western flank of the Sierras de Córdoba (Miranda and Introcaso, 1996) and is marked by the Guzman Fault in San Luis.

### 4.1 EARLY CAMBRIAN SEDIMENTATION

The oldest rocks in the region form a structurally thick sequence of paragneisses of the Pinchanas Metamorphic Complex. These metasediments are interpreted as being deposited on a passive margin, developed during intracontinental rifting and break up of Laurentia from Gondwana in Eocambrian times at about 540Ma (Dalziel and others 1994) in a tectonic environment similar to that envisaged by Dalla Salda and others (1994). Lithological similarities and comparable ages indicate that the metasediments may be correlatives of the Early Cambrian (Aceñolaza and Toselli, 1981) Puncoviscana

Formation in the northern Sierras Pampeanas as postulated by Willner and Miller (1986).

## 4.2 PAMPEAN CYCLE

### *Early Cambrian deformation, metamorphism, mafic and felsic intrusion*

Following intrusion of rare tholeiitic mafic dykes, the sediments were deformed at mid-crustal levels by a compressive event (D1) and metamorphosed at mostly upper amphibolite facies and locally, granulite-facies to form banded gneiss and locally migmatites. Mafic dykes were converted to amphibolite and extensively boudinaged. Muscovite-pegmatites formed subconcordant lenses. Estimates of peak metamorphic conditions for the area are mostly about 6 Kb, and 700°C to 800°C. A penetrative differentiated foliation formed as the last, and possibly second or even third fabric, in a progressive westerly-directed thrusting event, which is evident in the rocks by the presence of mylonite and a ubiquitous east-plunging mineral lineation, commonly sillimanite. Uranium-lead dating of zircon rims formed during this metamorphic event (M1) in Córdoba give an age of about 530 Ma (Camacho and Ireland, 1997). In the map region this event includes both the D1 and D2 domains of Dalla Salda (1987) and has been previously termed the “Ciclo orogénico Pampeano” (Aceñolaza and Toselli, 1976) or “Ciclo Pampeano” (Dalla Salda, 1987, Toselli and others, 1992). The deformation is interpreted as the first in a series of deformation events associated with convergence on the newly created Pacific Gondwana margin formed after final amalgamation of the supercontinent (e.g. Dalziel and others, 1994).

## 4.3 FAMATINIAN CYCLE

### *Early Ordovician Deformation, metamorphism, mafic and felsic intrusion*

During the Ordovician, closure of the Iapetus Ocean and collision of the Precordillera with the Pampean margin of the Gondwana craton (Dalla Salda and others, 1992, 1996, Dalziel and others, 1996) resulted in amalgamation of the accretionary wedge and the Pampean domain during a widespread deformational, metamorphic and magmatic event known as the “Ciclo orogénico Famatiniano” (Aceñolaza and Toselli, 1976), Famatinian Orogen (e.g. Dalla Salda and others, 1992) or “Ciclo Famatiniano” (Dalla Salda, 1987). Earlier D1 fabrics were tightly folded with local axial plane crenulation cleavage, subparallel to the higher grade differentiated D1 fabric.

Dalla Salda (1987) and Toselli and others (1992) ascribed this deformation to the D2 domain. Zircons which grew during this event yield an age of about 480 Ma dating the timing of peak metamorphism in the Famatinian domain in San Luis.

Compression was followed closely, possibly in the one continuous event, by extensional deformation (D2 in the Famatinian domain) at amphibolite- to greenschist-facies. This deformation was mostly confined to the already established ductile shear-zones and was accompanied by retrogression of higher grade metamorphic assemblages and by intrusion of numerous granitic to tonalitic bodies and tourmaline-bearing pegmatites. It is not known whether the extensional phase affected the Pichanas Metamorphic Complex in the 3166-23 Sheet area.

#### 4.4 ACHALIAN CYCLE

##### *Early Devonian granite intrusion and deformation*

Mid Palaeozoic resumption of convergence on the western margin of Gondwanaland is evidenced by a widespread compressive deformation in the Famatinian (D2) and Pampean domains (D3), and the development of an Early Devonian magmatic arc. The deformation was dominated by orthogonal westerly-directed thrusting and the development of regionally extensive ductile shear zones with intensive greenschist facies retrogressive fabrics. Locally, outside the principal shear zones, the metamorphic rocks were open to isoclinally folded with an axial crenulation developed in places.

Dalla Salda (1987) defined this deformation as D3, placing it in the “Ciclo Famatiniano”.

Peraluminous to slightly peralkaline felsic melts, generated from partial melting of MgO depleted crustal rocks (Dalla Salda and others, 1995) intruded the metamorphics discontinuously during and after shear zone development.). U-Pb zircon dating of these granites in the southern Sierras Pampeanas brackets crystallisation of the felsic magmas and shear zone formation over a 20 Ma period between 404 Ma and 384 Ma. The Achalian Cycle derives its name from the Achala Batholith, the largest of the Devonian batholiths in the southern Sierras Pampeanas, which is exposed south of the map area in the Sierra Grande. The cycle probably corresponds to the “Fase Precordilleránica” (Astini, 1996) in the precordillera west of the Sierras Pampeanas where it is related to the amalgamation of the Chilena domain.

The final stages of the Achalian Cycle were the province-wide development of a complex system of rectilinear brittle-ductile vertical NW- and NE-trending strike-slip faults and fractures. The orientation and conjugate relationship of the fractures indicates a continuation of the east-west compressive regime. Locally, the structures are associated with vein-type Au±Cu mineralisation, the result of mesothermal activity interpreted to be associated with the waning stages of magmatic arc activity as the centre of magmatic activity migrated westward (Ramos and others, 1986). Muscovite Ar-Ar ages indicate that this stage began about 385 Ma, peaked at 370 Ma and continued until 355 Ma (Camacho, 1997). Toselli and others (1996) attribute development of the fracture system to a 355 Ma old “Chánica Orogeny”.

#### **4.5 CARBONIFEROUS - PERMIAN SEDIMENTATION**

Following peneplanation, and later marine transgression, fluvio-lacustrine and shallow-marine sediments of the Paganzo Group (González and Aceñolaza, 1972) were deposited during the Carboniferous and Permian. The sediments may have covered much of the crystalline basement, however, only remnant outcrops of the group are now preserved in



narrow (<2 km wide) grabens. These grabens, possibly initiated during syn-sedimentary extensional faulting, were active after the cessation of sedimentation and prior to the Andean Cycle deformation. It is possible that the late-Palaeozoic sediments were first deposited in basins controlled by the regional wrench tectonic regime.

#### 4.6 ANDEAN CYCLE

East-west compression during the Cainozoic Andean uplift resulted in Neogene inversion of the Cretaceous basins (Schmidt, 1993) and block tilting of basement rocks, forming north-south oriented ranges separated by intermontane basins. The ranges are bounded by escarpments developed on moderate to steeply-dipping reverse faults (Jordan and Allmendinger, 1986; Martino and others, 1995; Costa, 1996), many of which show repeated reactivation. Costa interpreted the last and most significant movement in the region took place during the Late Pliocene-Pleistocene with some movement continuing during the Quaternary.

## SECTION 2: ECONOMIC GEOLOGY

by Roger G. Skirrow

### 1. INTRODUCTION

The northern 0°5' portion of Sheet 3166-23 covered in this report contains few metallic mineral occurrences, although a variety of commodities are represented including W, Ag-Pb-Zn, Fe and Be. Travertine is quarried near La Playa.

In the Geoscientific Mapping of Sierras Pampeanas Cooperative Project the principal metallic deposits in all main mining districts of the map area were investigated in the field, and geological observations were entered into the ARGROC and ARGMIN databases (Skirrow and Trudu, 1997). ARGMIN is a Microsoft Access database that was initially developed jointly by AGSO and the Subsecretaría de Minería in ORACLE, based on OZMIN (Ewers and Ryburn, 1993). Additional geological and resource data from the literature on mineral occurrences have been compiled in ARGMIN. Petrography of ore and host rock samples (thin sections and polished thin sections) was recorded in a petrographic database (Sims and others, 1996), and selected samples for ore genesis studies were analysed for whole rock geochemistry (Lyons and others, 1996; Lyons and Skirrow, 1996), stable isotopes of oxygen, hydrogen and sulfur (Lyons and Skirrow, 1986), as well as  $^{40}\text{Ar}/^{39}\text{Ar}$  radiometric age dating (Camacho, 1997). Geographic coordinates were measured by GPS (locational accuracy 50m), whereas those occurrences not visited in the field were generally located on aerial photographs and their geographic coordinates digitised. The locational accuracy for photo-located occurrences is 200 m. The locations of remaining occurrences are taken from various published sources, which in some cases allow only very approximate geographic coordinates to be estimated (e.g. 3 km for U deposits).

Mineral occurrence data as well as non-metallic mineral and dimension stone occurrences are shown on the 1:100 000 scale metallogenic map accompanying this report. Output data sheets from the ARGMIN database are appended to the report. Details of the geology and grade-tonnage data, where available, for individual metallic mineral occurrences may be found in the database. The 1:250 000 scale Metallogenic Map for the Sierras Septentrionales de Córdoba (Skirrow, 1997) shows the mineral occurrences in relation to 'prospectivity domains' or areas of mineral potential. These domains are defined on the basis of 'metallogenic models' for each mineral deposit style, as discussed by Skirrow Lyons and others (1997) and which were developed from the observations and interpretations presented in the following sections. For further datasets of mineral potential, the reader is referred to the *Atlas Metalogénico* (1:400 000 scale) for the Sierras Pampeanas mapping project (Skirrow and Johnston, 1997) and project GIS (Butrovski, 1997) in which metallogenic models for the principal styles of metallic mineralisation are presented as separate coverages.

## **2. METALLIC MINERAL OCCURRENCES**

Occurrences of W, Ag-Pb-Zn, Fe and Be in the northern 0°5' portion of Sheet 3166-23 are known only from the 1:750 000 scale map of Ricci (1974). No descriptions of the geology or accurate locations are available.

## **3. NON-METALLIC MINERALS AND *ROCAS DE APLICACION***

### **3.1 LIMESTONE, MARBLE**

Small bodies of travertine carbonate occur near La Playa (Lucero Michaut and Olascher, 1981). The travertine probably represents low temperature hydrothermal activity associated with Miocene-Pliocene volcanism in the Pocho region to the south.

### **3.2 GRANITE**

Dimension stone is obtained from the La Playa Granodiorite.

### **3.3 FLUORITE, CLAY, OCHRE, STEATITE, GARNET AND AMPHIBOLITE**

Several worked occurrences of fluorite, clay, ochres, steatite, garnet, amphibolite were shown on the maps of Ricci (1974), Pastore and Methol (1953) and Lucero Michaut and Olascher (1981), and on the 1995 Mapa Geológica de la Provincia de Córdoba (1:500 000 scale).

### **3.4 MICA, QUARTZ, FELDSPAR**

Numerous relatively small pegmatite bodies have been worked for muscovite, quartz and feldspar and occur widely throughout the region (Ricci, 1974).

## BIBLIOGRAPHY

- ACENOLAZA, F.G., y TOSELLI, A.J., 1976. Consideraciones estratigráficas y tectónicas sobre el Paleozoico inferior del Noroeste Argentino. Memoria, II Congreso Latinoamericano de Geología, 2, 755-764.
- ACENOLAZA, F.G., y TOSELLI, A.J., 1981. Geología de Noroeste Argentino. Publicación especial Fac. Ci. Nat. UNT, Tucumán, 1287, 212 p
- ASTINI, R.A., 1996. Las fases diastroficas del Paleozoico medio en La Precordillera del oeste Argentino - evidencias estratigráficas. XIII Congreso Geológico Argentino y III Congreso de Exploración de Hidrocarburos, Actas V: 509-526.
- BUTROVSKI, D., 1997. Geographic Information System (GIS) for the Sierras Pampeanas Mapping Project, Argentina. Australian Geological Survey Organisation, Arc/Info GIS.
- CAMACHO, A., 1997.  $^{40}\text{Ar}$ - $^{39}\text{Ar}$  and Rb-Sr geochronology, final report. Geoscientific mapping of the Sierras Pampeanas, Argentine-Australia Cooperative Project, Australian Geological Survey Organisation, unpublished report.
- CAMACHO, A. and IRELAND, T.R., 1997. U-Pb geochronology, final report. Geoscientific mapping of the Sierras Pampeanas, Argentine-Australia Cooperative Project, Australian Geological Survey Organisation, unpublished report.
- COSTA, C.H., 1996. Analisis neotectónico en las sierras de San Luis y Comechingones: problemas y methods. XIII Congreso Geológico Argentino y III Congreso de Exploración de Hidrocarburos, Actas II: 285-300.
- DALLA SALDA, L.H., 1987. Basement tectonics of the southern Pampean ranges, Argentina. *Tectonics*, 6: 249-260
- DALLA SALDA, L.H., CINGOLANI, C., VARELA, R., and LOPEZ DE LUCHI, M., 1995. The Famatinian Orogenic Belt in South-western South America: granites and metamorphism: an Appalachian similitude?. IX Congreso Latinoamericano de Geología, Caracas, Resúmenes.
- EWERS, G.R. and RYBURN, R.J., 1993. User's guide to the OZMIN mineral deposit database. Australian Geological Survey Organisation, Record 1993/94, 69p.
- FITZSIMMONS, I.C.W., KINNY, P.D., and HARLEY, S.L., 1996. Timescales of Pan-African metamorphism, melting and exhumation at Brattstrand Bluffs, east Antarctica. *Evolution of Metamorphic Belts*, Geological Society of Australia, Abstract volume 42.
- GONZALE, R.R. y ACENOLAZA, F.G. 1972. La cuenca de deposición neopaleozóica-mesozóica del oeste argentino. *Fundación e Instituto Miguel Lillo*, Tucumán, *Miscelánea*, 40: 629-643.
- JORDAN, T.E. AND ALLMENDINGER, R.W. 1986. The Sierras Pampeanas of Argentina: A modern analogue of Rocky Mountain foreland deformation. *American Journal of Science*, 286: 737-764.

- LAPIDUS, A.L. y ROSSI, N.I., 1959. Las minas de tungsteno de Agua de Ramón, Departamento Minas, provincia de Córdoba. Ministerio de Economía de la Nación, Anales 10. Buenos Aires.
- LUCERO MICHAUT, H.N. y OLASCHER, J., 1981. Descripción Geológica de la Hoja 19h, Cruz del Eje, Provincia de Córdoba (Escala 1:200 000). Servicio Geológico Nacional, Boletín, 179, 96 p.
- LYONS, P. and SKIRROW, R.G., 1996. Whole rock and stable isotope geochemistry - Final Report. Geoscientific Mapping of the Sierras Pampeanas Argentine-Australian Cooperative Project, Australian Geological Survey Organisation, unpublished report.
- LYONS, P, SKIRROW, R.G. and STUART-SMITH, P.G. 1997. Report on Geology and Metallogeny of the Sierras septentrionales de Córdoba, 1:250 000 map sheet, Province of Córdoba. Geoscientific Mapping of the Sierras Pampeanas, Argentina-Australia Cooperative Project, Australian Geological Survey Organisation, unpublished report.
- LYONS, P., STUART-SMITH, P.G., SIMS, J.P., PIETERS, P., SKIRROW, R.G. and CAMACHO, A., 1996. Whole Rock Geochemistry Report. Geoscientific Mapping of the Sierras Pampeanas Argentine-Australian Cooperative Project, Australian Geological Survey Organisation, unpublished report, June 1996.
- MARTINO, R., KRAEMER, P., ESCAYOLA, M., GIAMBASTIANI, M., and ARNOSIO, M., 1995. Transect de Las Sierras Pampeanas de Córdoba a los 32° S. Revista de la Asociación Geológica Argentina, 50: 60-77.
- MIRANDA, S. y INTROCASO, A., 1996. Cartas gravimétricas y comportamiento isostático areal de las Sierras de Córdoba - Rep. Argentina. XIII Congreso Geológico Argentino y III Congreso de Exploración de Hidrocarburos, Actas II: 405-417.
- OLSACHER, J., 1960. Descripción geológica de la hoja 20h, Los Gigantes, Provincia de Córdoba. Dirección Nacional de Geología y Minería. Buenos Aires. Bull. 90.
- PASTORE, F., 1932. Descripción de la Hoja Geológica 20I, Córdoba. Ministerio Agricultura, Dirección de Minería y Geología, Boletín, 36, 67pp.
- PASTORE, F. y METHOL, E. J., 1959. Las minas de tungsteno de Aguas de Ramón. Anales de Ministerio de Economía de la Nación,
- RAMOS, V.A., JORDAN, T.E., ALLMENDINGER, R.W., MPODOZIS, C., KAY, S., CORTES, J.M., and PALMA, M.A., 1986. Paleozoic terranes of the Central Argentine-Chilean Andes. Tectonics, 5: 855-880.
- RAPELA, C.W., PANKHURST, R.J., BALDO, E., and SAAVEDRA, J., 1995. Cordierites in S-type granites: Restites following low pressure, high degree partial melting of metapelites. The Origin of Granites and Related Rocks, Third Hutton Symposium Abstracts. US Geological Survey Circular 1129.
- RAPELA, C.W., SAAVEDRA, J., TOSELLI, A., AND PELLITERO, E, 1996. Eventos magmáticos fuertemente peraluminosos en las Sierras Pampeanas. XIII Congreso Geológico Argentino y III Congreso de Exploración de Hidrocarburos, Actas V: 337-353

- RICCI, S.M., 1974. Provincia de Córdoba - Mapa Minero, Escala 1 : 750,000, Ministerio de Industria y Minería, Subsecretaría de Minería, Dirección Nacional de Promoción Minera.
- SANTA CRUZ, J.N., 1978. Aspectos sedimentológicos de las formaciones aflorantes al este de la Sierra Chica, Provincia de Córdoba, Republica Argentina. Asociación Geológica Argentina, revista, 36: 232-244.
- SCHMIDT, C., 1993. Neogene inversion of two Cretaceous basins, Sierras Pampeanas, Argentina. Geological Society of America, 1993 Annual Meeting, Boston, Abstracts, 233.
- SIMS, J.P., STUART-SMITH, P.G, LYONS, P. y SKIRROW, R., 1997. Sierras de San Luis y Comechigones (Informe 1:250.000 Hoja geología). Servicio Geológico Minero Argentino - Australian Geological Survey Organisation, unpublished report.
- SIMS, J.P., STUART-SMITH, P.G., LYONS, P., PIETERS, P., SKIRROW, R.G. and CAMACHO, A., 1996. Petrography Report. Geoscientific Mapping of the Sierras Pampeanas Argentine-Australian Cooperative Project, Australian Geological Survey Organisation, unpublished report, June 1996.
- SKIRROW, R.G. y JOHNSTON, A.I., 1997. Atlas Metalogénico de las Sierras Pampeanas, República Argentina. Australian Geological Survey Organisation, Geoscientific Mapping of the Sierras Pampeanas Argentine-Australian Cooperative Project.
- SKIRROW, R.G. and TRUDU, A., 1997. ARGMIN: a mineral deposit database for the Sierras Pampeanas, Republic of Argentina. Australian Geological Survey Organisation, Geoscientific Mapping of the Sierras Pampeanas Argentine-Australian Cooperative Project. Database in Microsoft Access and Oracle.
- STRASSER, E.N., TOGNELLI, G.C., CHIESA, J.O. y PRADO, J.L., 1996. Estratigrafía y sedimentología de los depósitos eólicos del pleistoceno tardío y Holoceno en el sector sur de la sierra de San Luis. XIII Congreso Geológico Argentino y III Congreso de Exploración de Hidrocarburos, Actas IV: 73-83.
- TOSELLI, A.J., DALLA SALDA, L. y CAMINOS, R., 1992. Evolución metamórfica del Paleozoico Inferior de Argentina. In J.G. Gutiérrez Marco, J Saavedra and I. Rábano (Eds), Paleozoico Inferior de Ibero-América. Universidad de Extremadura.
- TOSELLI, A.J., DURAND, F.R., ROSSI DE TOSELLI, J.N. y SAAVEDRA, J., 1996. Esquema de evolución geotectónica y magmática eopaleozoica del Sistema de Famatina y sectores de Sierras Pampeanas. XIII Congreso Geológico Argentino y III Congreso de Exploración de Hidrocarburos, Actas V: 443-462.
- WILLNER, A.P., and MILLER, H., 1986. Structural division and evolution of the lower Paleozoic basement in the NW Argentine Andes. Zentralblatt für Geologie und Paläontologie, I, 1245-1255.
- ZOLEZZI, R.E., MIRÓ, R.C. y PETRELLI, H., 1988. Perfil del Proyecto Puigari-Monserrat, Secretaría de Minería, Dirección Nacional de Minería y Geología, Centro de Exploración Córdoba, 19 p.

**ARGMIN**  
**DATABASE OUTPUT SHEETS**